

correction dose

This is the insulin dose you take for high blood glucose during the day and at bedtime.

Meal time: 1 unit per 10mg/dL over 100

Bedtime: 1 unit per 10mg/dL over 110

blood glucose	daytime	bedtime
Under 100	0	0
100-110	1	0
111-120	2	1
121-130	3	2
131-140	4	3
141-150	5	4
151-160	6	5
161-170	7	6
171-180	8	7
181-190	9	8
191-200	10	9
201-210	11	10
211-220	12	11
221-230	13	12
231-240	14	13
241-250	15	14
251-260	16	15

continued on back

Only give correction dose if it has been **more than 2-3 hours** since the last dose of fast-acting insulin.

blood glucose	daytime	bedtime
261-270	17	16
271-280	18	17
281-290	19	18
291-300	20	19
301-310	21	20
311-320	22	21
321-330	23	22
331-340	24	23
341-350	25	24
351-360	26	25
361-370	27	26
371-380	28	27
381-390	29	28
391-400	30	29
401-410	31	30
411-420	32	31
421-430	33	32
431-440	34	33
441-450	35	34
451-460	36	35
461-470	37	36
471-480	38	37
481-490	39	38
491-500	40	39
Over 500	41	40

childrensdayton.org/diabetesresources

