

correction dose

This is the insulin dose you take for high blood glucose during the day and at bedtime.

Meal time: 1 unit per 5mg/dL over 125

Bedtime: 1 unit per 5mg/dL over 130

blood glucose	daytime	bedtime
Under 125	0	0
125-130	1	0
131-135	2	1
136-140	3	2
141-145	4	3
146-150	5	4
151-155	6	5
156-160	7	6
161-165	8	7
166-170	9	8
171-175	10	9
176-180	11	10
181-185	12	11
186-190	13	12
191-195	14	13
196-200	15	14
201-205	16	15
<i>continued on back</i>		

Only give correction dose if it has been **more than 2-3 hours** since the last dose of fast-acting insulin.

blood glucose	daytime	bedtime
206-210	17	16
211-215	18	17
216-220	19	18
221-225	20	19
226-230	21	20
231-235	22	21
236-240	23	22
241-245	24	23
246-250	25	24
251-255	26	25
256-260	27	26
261-265	28	27
266-270	29	28
271-275	30	29
276-280	31	30
281-285	32	31
286-290	33	32
291-295	34	33
296-300	35	34
301-305	36	35
306-310	37	36
311-315	38	37
316-320	39	38
321-325	40	39
326-330	41	40
331-335	42	41

childrensdayton.org/diabetesresources

